

Quantum-Fluctuation-Driven Coherent Spin Dynamics in Small Condensates

Xiaoling Cui^{1,2}, Yupeng Wang¹ and Fei Zhou²

¹*Beijing National Laboratory for Condensed Matter Physics and Institute of Physics,
Chinese Academy of Sciences, P. O. Box 603, Beijing 100190, China*

²*Department of Physics and Astronomy, The University of British Columbia, Vancouver, B. C., Canada V6T1Z1*

(Dated: June 6, 2008)

We have studied quantum spin dynamics of small condensates of cold sodium atoms. For a condensate initially prepared in a mean field ground state, we show that coherent spin dynamics are *purely* driven by quantum fluctuations of collective spin coordinates and can be tuned by quadratic Zeeman coupling and magnetization. These dynamics in small condensates can be probed in a high-finesse optical cavity where temporal behaviors of excitation spectra of a coupled condensate-photon system reveal the time evolution of populations of atoms at different hyperfine spin states.

Recently, single-atom detection in optical cavities has been realized in experiments by having atoms and cavity photons in a strongly coupling regime[1, 2]. This remarkable achievement has been applied to study optically transported atoms in cavities[3]; furthermore the coupling between a small Bose-Einstein condensate (BEC) and cavity photons and resultant collective excitations have also been successfully investigated[4]. The sensitivity that a cavity-based atom detector has, together with a translating optical lattice which can effectively transport ultra cold atoms from a magnetic-optical trap to a cavity make it possible to study the physics of small BECs. Especially, this potentially opens the door to explore coherent dynamics of ultra-cold atoms in relatively small condensates. The physics of BECs of small numbers of atoms can be qualitatively different from the physics of big condensates and represents a new domain of cold-atom research. In small condensates, various intrinsic beyond-mean-field dynamics can be relevant within an experimentally accessible time scale. These new physical phenomena however have been quite difficult to study using the standard absorption-imaging approach to cold atoms because of relatively fewer atoms are involved in small condensates. Cavity electrodynamics in a strong coupling regime and high sensitivities to intra-cavity atoms on the other hand are ideal for investigating small condensates where the beyond-mean-field dynamics are mostly visible. In this letter, we focus on the basic concepts of beyond-mean-field coherent spin dynamics in BECs with typically a few tens to a few hundreds of atoms and detailed analysis of detecting these fascinating properties of small condensates in optical cavities with high-finesse. Research on this subject could substantially advance our understanding of the nature of quantum-fluctuation dynamics[5], in this particular case, dynamics purely driven by fluctuations with wavelengths of the size of condensates. Secondly, results obtained can help to better recognize limitations of precise measurements of various interaction constants based on mean-field coherent dynamics[6]. Thirdly, our results should shed some light on the feasibility of investigating fluctuation dynamics of small condensates using optical

cavities and also pave the way for future studies of dynamics of coupled small condensates.

To understand spin dynamics of a small condensate, we first study the evolution of a condensate of N hyperfine spin-one sodium atoms which is initially prepared in a mean field ground state,

$$|\mathbf{n}\rangle = \frac{(\mathbf{n} \cdot \boldsymbol{\psi}^\dagger)^N}{\sqrt{N!}} |0\rangle. \quad (1)$$

Here $\mathbf{n}$ is a unit director and three components of $\boldsymbol{\psi}^\dagger$, $\psi_\alpha^\dagger$, $\alpha = x, y, z$ are creation operators for three spin-one states, $|x\rangle = (|1\rangle - |-1\rangle)/\sqrt{2}$, $|y\rangle = (|1\rangle + |-1\rangle)/i\sqrt{2}$ and $|z\rangle = |0\rangle$ respectively. And in this representation, $S_\alpha = -i\epsilon_{\alpha\beta\gamma}\psi_\beta^\dagger\psi_\gamma$ is the total spin operator. States in Eq.1 with $\mathbf{n} = \mathbf{e}_z$ minimize the interaction energy of the following Hamiltonian for spin-one atoms in the presence of a quadratic Zeeman coupling along the z -direction,

$$H = \frac{c_2}{N} \mathbf{S}^2 + q(\psi_x^\dagger\psi_x + \psi_y^\dagger\psi_y). \quad (2)$$

Here c_2 is a spin interaction constant and q is the quadratic Zeeman coupling[7, 8, 9, 10]. Mean field ground states are stationary solutions to the multi-component Gross-Pitaevskii equations for spin-one atoms and dynamics of these initial states demonstrated below are therefore a beyond-mean field phenomenon. When deriving Eq.2 for a trapped condensate, we assume that spin dynamics are described by a single mode, i.e. $\psi_\alpha(\mathbf{r}, t) = \sqrt{\rho(\mathbf{r})}\psi_\alpha(t)$; for a small condensate of less than one thousand weakly interacting atoms, this approximation is always valid. c_2 is typically a few nano kelvin for sodium atoms; $q = (\mu_B B)^2/(4\Delta_{hf})$ and the hyperfine splitting is $\Delta_{hf} = (2\pi)1.77GHz$ (μ_B is the Bohr magneton and $\hbar$ is set to be unity).

To illustrate the nature of non-mean-field dynamics and crucial role played by quantum fluctuations, we expand the full Hamiltonian in Eq.2 around a mean field ground state. In the lowest order expansion, we approximate $\boldsymbol{\psi}^\dagger \approx \sqrt{N}\mathbf{e}_z + \psi_x^\dagger\mathbf{e}_x + \psi_y^\dagger\mathbf{e}_y$, and $\psi_{x,y}^\dagger$ are much

less than $\sqrt{N}$; the Hamiltonian then can be expressed in terms of the bilinear terms

$$H_B = \sum_{\alpha=x,y} \frac{q+4c_2}{2N} P_\alpha^2 + \frac{qN}{2} \theta_\alpha^2 + \dots \quad (3)$$

where for $\alpha = x, y$, $\theta_\alpha = \frac{1}{\sqrt{2N}}(\psi_\alpha^\dagger + \psi_\alpha)$ and $P_\alpha = i\sqrt{\frac{N}{2}}(\psi_\alpha^\dagger - \psi_\alpha)$ are pairs of conjugate operators which satisfy the usual commutation relations $[\theta_\alpha, P_\beta] = i\delta_{\alpha,\beta}$. Semiclassically, collective coordinates θ_α , $\alpha = x, y$ represent projections of $\psi^\dagger$ or order parameter $\mathbf{n}$ in the xy plane, and $P_{y(x)} \sim S_{y(x)}$ is the spin projection along the $y(x)$ -direction. The bilinear Hamiltonian is equivalent to a harmonic oscillator moving along the direction of $\theta_{x,y}$ with a mass $m_{eff} = \frac{N}{q+4c_2}$, a harmonic oscillator frequency $\omega = \sqrt{q(q+4c_2)}$ and effective spring constant qN ; the mass at $q = 0$ is induced by scattering between atoms. The excitation spectrum is $E_n = (n + 1/2)\omega$, $n = 0, 1, 2, \dots$. When $q = 0$, the Hamiltonian describes a particle moving in a free space.




FIG. 1: a) (color online) Time evolution of $N_0(t)$, the atom population at $|1, 0\rangle$ state for different quadratic Zeeman coupling q . Initially, all $N = 200$ atoms occupy $|1, 0\rangle$ state which corresponds to a mean field ground state. The inset is t_c , the time for the first dip in the $q = 0$ data, as a function of the number of atoms N . b) Time evolution of $N_0(t)$ for different magnetization m (here $q = 0$). All initial states are again mean-field ground states for given m . Inset is for $m = 0.98$. In this and Fig.2,3, $c_2 = (2\pi)50Hz$.

In the ground state of the bilinear Hamiltonian of Eq.3, $\langle\theta_\alpha\rangle = \langle P_\alpha\rangle = 0$ and $\mathbf{n}$ and $\langle\mathbf{S}\rangle$ have no projections in the xy plane. However, quantum fluctuations of

$\theta_{x,y}$ -coordinates in the ground state can be estimated as $\langle\theta_\alpha\theta_\alpha\rangle = \frac{1}{2N}\sqrt{\frac{q+4c_2}{q}}$. This is a measure of how strongly $\mathbf{n}$ fluctuates in the xy -plane. As expected, these quantum fluctuations are substantial only when q is small and are suppressed by a quadratic Zeeman field which effectively pins the order parameter along the z -direction. A direct calculation also shows that the amplitude of quantum fluctuations $\langle\theta_\alpha^2\rangle_{MF}$ in the *mean field ground state* defined in Eq.1 is $1/2N$. This indicates that the mean field ground state is a good approximation only when $q \gg 4c_2$. On the other hand, as q decreases and the effective spring constant gets smaller, the deviation becomes more and more severe. When q approaches zero, quantum fluctuations θ_α in the harmonic oscillator ground state become divergent implying that the mean field ground state is no longer a good approximation.

Indeed, the the energy of mean field ground state is $E_{MF} = \frac{q}{2} + c_2$ which is much higher than $\frac{1}{2}\omega$ when $q \ll c_2$; such a state corresponds to a highly excited wave packet, because of a relatively narrow spread along θ_α -directions and consequently an enormous kinetic energy associated with momenta P_α . We therefore expect that dynamics in this limit could dramatically differ from a stationary solution. Since the total number of atoms N is equal to $\sum_\alpha \psi_\alpha^\dagger \psi_\alpha$, the population of atoms at $|z\rangle$ (or $|1, 0\rangle$) state $N_0 = \langle\psi_z^\dagger \psi_z\rangle$ is directly related to quantum fluctuations of θ_α and P_α ,

$$N_0 = N + 1 - \sum_\alpha \left(\frac{N}{2} \langle\theta_\alpha^2\rangle + \frac{1}{2N} \langle P_\alpha^2\rangle \right). \quad (4)$$

Eq.4 shows that the time evolution of $N_0(t)$ is effectively driven by quantum fluctuations in θ_α and P_α ; a study of $N_0(t)$ probes underlying quantum- fluctuation dynamics.

For an initial state prepared in a mean field ground state with $\mathbf{n} = \mathbf{e}_z$ where all atoms condense in $|1, 0\rangle$ state, one finds that $\langle\theta_\alpha^2\rangle = \frac{1}{2N}$ and $\langle P_\alpha^2\rangle = \frac{N}{2}$. The evolution of such a symmetric Gaussian wave packet subject to the bilinear Hamiltonian can be solved exactly using the standard theory for harmonic oscillators. The wave packet will remain to be a Gaussian one with the width oscillating as a function of time. Qualitatively, because of the symmetry, only harmonic states with even-parity are involved in dynamics and therefore the oscillation frequency is 2ω . Furthermore during oscillations, the kinetic energy stored in initial wave packets is converted into the potential one and *vice versa*. Especially when $q \ll c_2$, oscillations are driven by the enormous initial kinetic energy associated with P_α ; the oscillation amplitude can be estimated by equating the total energy E_{MF} to the potential energy which leads to $\langle\theta_\alpha^2\rangle \sim c_2/(Nq)$. A straightforward calculation yields the time dependence of $\langle\theta_\alpha^2\rangle$ and $\langle P_\alpha^2\rangle$ that leads to

$$\frac{N_0}{N} = 1 - \frac{8c_2^2}{q(q + 4c_2)N} \sin^2 wt. \quad (5)$$

The oscillating term in Eq.5 shows the deviation from the stationary behavior due to quantum fluctuations in θ_α -coordinates. The deviation is insignificant when q is not too small; however when q is of the order of c_2/N , we expect that the non-mean field dynamics becomes very visible. Note that the approach outlined here neglects all higher order anharmonic interactions and therefore is only valid when the relative amplitude of fluctuations is small; that is when $q \gg c_2/N$.

When q approaches zero, the short time dynamics following the bilinear Hamiltonian is equivalent to a particle of a mass $m_{eff} = N/4c_2$ that is initially localized within a spread $\langle \theta_\alpha^2 \rangle = 1/2N$ having a ballistic expansion with a typical velocity give as $\langle v_\alpha^2 \rangle = 8c_2^2/N$. The time dependence of spread $\langle \theta_\alpha^2 \rangle$ therefore is $1/2N + (8c_2^2/N)t^2$. So at $t \sim \sqrt{N}/c_2$, the number of atoms not occupying the initially prepared $|1,0\rangle$ state becomes of order of N . This limit was first addressed by Law *et al.* in the context of four-wave-mixing theory[9], and also in early works[11, 12]; to describe the physics after this characteristic time scale requires analysis of full quantum dynamics. This time scale however becomes quite long for a few million atoms which makes it difficult to observe quantum dynamics in large condensates.

In the following, we are going to present our numerical results on dynamics and focus on its dependence on quadratic Zeeman coupling q and magnetization m . For a condensate of $N = 200$ atoms, we numerically integrate the time-dependent N-body Schrodinger equation of the quantum Hamiltonian in Eq.2. The time evolution of N_0 driven by quantum fluctuations is shown in Fig.1a). As q increases far beyond $0.2c_2$, N_0 oscillates as a function of time with frequency 2ω and the amplitude of oscillations decreases; the damping is not visible over tens of oscillations. When q is below $0.2c_2$, anharmonic effects become substantial and oscillations are no longer perfect; when $q = c_2/40$, oscillations are strongly damped after a few cycles and revived afterwards. For $q = 0$, N_0 drops to a minimum of about $0.38N$ when $t = t_c = 0.53\sqrt{N}/c_2$ and remains to be a constant before reviving to be $0.8N$ at about $10t_c$. For sodium atoms with a typically density $2 \times 10^{14} \text{ cm}^{-3}$, $c_2 = (2\pi)50 \text{ Hz}$; $t_c = 23.8 \text{ ms}$ for $N = 200$ and increases to a few seconds when N reaches 2×10^6 .

We have also studied the quantum dynamics of a mean field condensate with a finite magnetization along the z -direction, $\mathbf{m} = m\mathbf{e}_z$. States which minimize the mean field energy of the Hamiltonian in Eq.2 with $q = 0$ are

$$|m\rangle = \frac{[(\cos \eta \mathbf{e}_x + i \sin \eta \mathbf{e}_y) \cdot \psi^\dagger]^N}{\sqrt{N!}} |0\rangle \quad (6)$$

where $\sin 2\eta = m$, $m(\in [-1, 1])$ is the normalized magnetization. By expanding the Hamiltonian around these mean field states, one obtains a harmonic oscillator Hamiltonian defined in terms of conjugate operators, $\theta_z = \frac{1}{\sqrt{2N}}(\psi_z^\dagger + \psi_z)$ and $P_z = i\sqrt{\frac{N}{2}}(\psi_z^\dagger - \psi_z)$. The effective mass is $m_{eff} = \frac{N}{2c_2(1+\sqrt{1-m^2})}$ and the harmonic oscillator frequency $\omega = 2|m|c_2$. States shown in Eq.6 have a narrow width along the direction of θ_z , $\langle \theta_z^2 \rangle = 1/2N$ and therefore carry large conjugate momenta P_z ; the corresponding large kinetic energy drives a unique non-mean field quantum spin dynamics. The harmonic expansion again is only valid when m is large and fluctuations are weak. Simulations of the full Hamiltonian have been carried out in this case; in Fig.1b), we show the time dependence of $N_0(t)$ for different magnetization. Only when m is close to unity, weakly damped oscillations are observed.




FIG. 2: (a,b,c) (color online) Eigenfrequencies Δ_p as a function of t for different detuning Δ_c when the relative population at state $|1,0\rangle$, $\rho_0(t)$ evolves. At $t = 0$, all atoms occupy $|1,0\rangle$ state and $N = 200$. d) Δ_p as a function of time t for $\Delta_c = 0$.

We propose a method to probe quantum spin dynamics of a small condensate of spin-one sodium atoms using cavity quantum electrodynamics. For a Bose-Einstein condensate with N atoms coupled to a quantized field of a cavity, a single cavity photon can coherently interact with atoms which leads to a collective coupling of $g\sqrt{N}$ [13]. In experiments[3, 4], atoms are transported into a cavity via a moving optical lattice; excitations are measured by individual recordings of cavity transmission when frequencies of an external probe light are scanned. Here we consider a multi-component BEC coupled to a single cavity mode; the eigenfrequencies of the coupled system uniquely depend on populations at three hyperfine states. By measuring the energy spectrum of this coupled system, one obtains temporal behaviors of atom

populations at different states.

We restrict ourselves to excitations which involve a single cavity photon interacting with atoms in a BEC. We study atomic transitions from $3S_{\frac{1}{2}} \rightarrow 3P_{\frac{1}{2}}$ in sodium atoms. The Hamiltonian consists the following terms,

$$H_{\text{cavity}} = \sum_i \hbar w_{g_i} \hat{g}_i^\dagger \hat{g}_i + \sum_j \hbar w_{e_j} \hat{e}_j^\dagger \hat{e}_j + \sum_p \hbar w_c \hat{c}_p^\dagger \hat{c}_p - i\hbar \sum_p \sum_{i,j} g_{ij}^p \hat{e}_j^\dagger \hat{c}_p \hat{g}_i + h.c., \quad (7)$$

where i labels three states $|F=1, m_F=0, \pm 1\rangle$ in $3S_{\frac{1}{2}}$ orbital and j eight states $|F'=1, m_{F'}\rangle, |F'=2, m_{F'}\rangle$ in $3P_{\frac{1}{2}}$ orbital. $\hat{g}_i^\dagger$ and $\hat{e}_j^\dagger$ create atoms in one of $3S_{\frac{1}{2}}$ and $3P_{\frac{1}{2}}$ states respectively with corresponding frequencies w_{g_i}, w_{e_j} . $\hat{c}_p^\dagger$ creates a photon with frequency w_c and polarization p in the cavity mode. $g_{ij}^p (= D_{ij}^p \sqrt{\hbar w_c / 2\epsilon_0 V})$ is the coupling strength for a transition $i \rightarrow j$ driven by a cavity photon with polarization p , which depends on the dipole matrix element D_{ij}^p , the effective mode volume V .




FIG. 3: Eigenfrequencies Δ_p as a function of time t driven by dynamics of population $\rho_0(t)$ in the presence of quadratic Zeeman coupling $q = 0.05c_2$ (or $B = 95mG$). Again at $t = 0$, all atoms occupy $|1, 0\rangle$ state and $N = 200$, $\Delta_c = 0$.

For simplicity, we set the energy of $3S_{\frac{1}{2}}$ states to be zero, i.e. $w_{g_i} = 0$; the energy of excited states is $w_e^{1,2} = w_a \pm \Delta$ with 2Δ being the hyperfine splitting between $F' = 2$ and $F' = 1$ states. For atomic transitions induced by left-circularly (σ^+) polarized photons, the selection rule is $\Delta F = 0, \pm 1, \Delta m_F = 1$. In a cavity, a state with a cavity photon (labeled as $1c$), N_{m_F} atoms at $3S_{\frac{1}{2}} |1, m_F\rangle$ states and no atoms at excited states (labeled as 0_j) is expressed as $|1c; N_1, N_0, N_{-1}; 0_j\rangle$; it is coupled to the following states with one of atoms excited to $3P_{\frac{1}{2}}$ states (labeled $1_{F'}$) and no cavity photons (as 0_c), $|0_c; N_1 - 1, N_0, N_{-1}; 1_{F'=2}\rangle$, $|0_c; N_1, N_0 - 1, N_{-1}; 1_{F'=1}\rangle$, $|0_c; N_1, N_0 - 1, N_{-1}; 1_{F'=2}\rangle$, $|0_c; N_1, N_0, N_{-1} - 1; 1_{F'=1}\rangle$, $|0_c; N_1, N_0, N_{-1} - 1; 1_{F'=2}\rangle$. We diagonalize the Hamiltonian matrix and obtain six eigenfrequencies ω_p for this coupled system. Three are $3P_{\frac{1}{2}}$ orbitals without mixing with $3S_{\frac{1}{2}}$ states, $w_p = w_a \pm \Delta$; the other three depend on *relative* populations of atoms at each spin state,

$\rho_{m_F} = N_{m_F}/N$, $m_F = 0, \pm 1$. The latter three eigenfrequencies are determined by the eigen value equation,

$$(\Delta_p - \Delta_c)(\Delta_p^2 - \Delta^2) - \Delta_p N g_1^2 F_1 - \Delta N g_1^2 F_2 = 0. \quad (8)$$

Here $\Delta_p = w_p - w_a$, $m = \rho_{+1} - \rho_{-1}$ is the normalized magnetization, and g_1 is the coupling between $3S_{\frac{1}{2}} |F=1, m_F=1\rangle$ and $3P_{\frac{1}{2}} |F'=2, m_{F'}=2\rangle$ by σ^+ light; $F_1 = (2+m)/3$ and $F_2 = (1+m)/2 - \rho_0/6$. Apparently eigenfrequencies $\Delta_{p1,p2,p3}$ are a function of ρ_{m_F} and therefore can be used to probe the variation in $\rho_{0,\pm 1}$ due to coherent spin dynamics. $\Delta_{p1,p2,p3}$ depend on a dimensionless parameter $r = \frac{\sqrt{6}}{3} \frac{\sqrt{N} g_1}{\Delta}$. When detuning $\Delta_c = 0$ and as $r \rightarrow \infty$, $\Delta_{p1,p2,p3}$ are around $0, \pm \frac{\sqrt{6}}{3} \sqrt{N} g_1$ respectively. Eigenfrequency Δ_{p2} varies from $-\Delta/4$ to $-\Delta/2$ when ρ_0 increases from 0 to 1; the variation amplitude $\delta = \Delta_p(\rho_0 = 1) - \Delta_p(\rho_0 = 0)$ reaches a saturated value $\Delta/4$. For sodium atoms, $\Delta = (2\pi)94.4MHz$; cavity parameters are chosen according to Ref.[4] and $g_1 = (2\pi)10MHz$. In Fig. (2), we show the evolution of Δ_p in time for different detuning Δ_c when atoms are initially prepared at state $|1, 0\rangle$ of $3S_{\frac{1}{2}}$. The evolution of $\Delta_p(t)$ which can be probed by a σ^+ beam maps out population $N_0(t)$ driven by underlying quantum fluctuations. In Fig. (3), we further show the time dependence of Δ_p due to oscillatory quantum spin dynamics for $q = 0.05c_2$ ($B \approx 95mG$), $N = 200$ and $\Delta_c = 0$.

In conclusion, we have illustrated the nature of coherent spin dynamics driven by quantum-fluctuations in small condensates. The time evolution of population of atoms at different hyperfine spin states is shown to reveal intrinsic dynamics of quantum fluctuations of order parameters and spin projections. These dynamics can be probed by studying eigenfrequencies of a coupled condensate-photon system in a high-finesse optical cavity available in laboratories. We thank Gerard Milburn, Junliang Song for stimulating discussions. This work is in part supported by NSFC, 973-Project (China), and NSERC (Canada), CIFAR and A. P. Sloan foundation.

-
- [1] H. Mabuchi, Q. A. Turchette, M. S. Chapman and H. J. Kimble, *Optics Letters* **21**, 1393 (1996).
 - [2] C. J. Hood *et al.*, *Phys. Rev. Lett.* **80**, 4157 (1998).
 - [3] J. A. Sauer *et al.*, *Phys. Rev. A* **69**, 051804(2004).
 - [4] F. Brennecke *et al.*, *Nature* **450**, 268 (2007).
 - [5] Other fluctuation-driven spin dynamics were studied in J. L. Song, F. Zhou, *Phys. Rev. A* **77**, 033628 (2008).
 - [6] M. S. Chang *et al.*, *Nature Physics* **1**, 111 (2005).
 - [7] T. L. Ho, *Phys. Rev. Lett.* **81**, 742 (1998).
 - [8] T. Ohmi, K. Machinda, *J.Phys.Soc.Jpn.* **67**, 1822 (1998).
 - [9] C. K. Law *et al.*, *Phys. Rev. Lett.* **81**, 5257 (1998).
 - [10] J. Stenger *et al.*, *Nature* **396**, 345(1998)
 - [11] F. Zhou, *Phys. Rev. Lett.* **87**, 080401 (2001).
 - [12] R. B. Diener and J. L. Ho, *cond-mat/0608732*(2006).
 - [13] M. Tavis, F. W. Cummings, *Phys. Rev.* **170**, 379 (1968).